ELSEVIER

Carbohydrate Research 281 (1996) C19–C21

CARBOHYDRATE RESEARCH

Contents

Regular papers

Crystal structure and solid-state NMR analysis of methyl 2,3,6-tri-*O*-acetyl-2-deoxy-2-[3-(2-phenylethyl)-ureido]-β-D-glucopyranoside
R. Anulewicz, I. Wawer, B. Piekarska-Bartoszewicz and A. Temeriusz (Warsaw, Poland) 1

Molecular dynamics simulations of maltose in water
K.-H. Ott and B. Meyer (Athens, GA, USA) 11

NMR determination of the structure of Julibroside J_1
L. Ma, G. Tu, S. Chen, R. Zhang, L. Lai, X. Xu and Y. Tang (Beijing, People's Republic of China). 35

Synthesis of the methyl α-glycosides of a di-, tri-, and a tetra-saccharide fragment mimicking the terminus of the O-polysaccharide of *Vibrio cholerae* O:1, serotype Ogawa
P.-s. Lei, Y. Ogawa and P. Kováč (Bethesda, MD, USA) 47

A convenient new pathway for stereospecific epimerization of monosaccharide moieties in disaccharides
R. Miethchen, D. Rentsch, M. Frank (Rostock, Germany) and A. Lipták (Debrecen, Hungary). . . . 61

Wittig chain extension of unprotected carbohydrates: formation of carbohydrate-derived α,β-unsaturated esters
C.J. Railton and D.L.J. Clive (Edmonton, Canada) 69

The aqueous solution structure of a lipoteichoic acid from *Streptococcus pneumoniae* strain R6 containing 2,4-diamino-2,4,6-trideoxy-galactose: evidence for conformational mobility of the galactopyranose ring
R.A. Klein, R. Hartmann, H. Egge (Bonn, Germany), T. Behr and W. Fischer (Erlangen-Nürnberg, Germany) 79

Synthesis of sulfur-linked analogues of nigerose, laminarabiose, laminaratriose, gentiobiose, gentiotriose, and laminaran trisaccharide Y
M.-O. Contour-Galcera, J.-M. Guillot, C. Ortiz-Mellet (Grenoble, France), F. Pflieger-Carrara (Aubière, France), J. Defaye (Grenoble, France) and J. Gelas (Aubière, France) 99

Stereocontrolled synthesis of sulfur-linked analogues of the branched tetrasaccharide repeating-unit of the immunostimulant polysaccharide schizophyllan and of its β-(1 → 3)-branched, β-(1 → 6)-linked isomer
M.-O. Contour-Galcera, Y. Ding, C. Ortiz-Mellet and J. Defaye (Grenoble, France) 119

Preparation of methyl 4,6-di-*O*-acetyl-3-*C*-nitro-2-enopyranoside derivatives and their reduction with sodium borodeuteride
A. Seta, K. Tokuda, M. Kaiwa and T. Sakakibara (Yokohama, Japan). 129

Structural characterization of extracellular polysaccharides produced by fungi from the genus *Pleurotus*
A. Gutiérrez, A. Prieto and A.T. Martínez (Madrid, Spain) 143

Structural studies of the *Escherichia coli* O26 O-antigen polysaccharide
M.C. Manca (Stockholm, Sweden), A. Weintraub (Huddinge, Sweden) and G. Widmalm (Stockholm, Sweden) 155

Structural features of an arabinan fragment isolated from the water-soluble fraction of dehulled rapeseed
I. Eriksson, R. Andersson, E. Westerlund, R. Andersson and P. Åman (Uppsala, Sweden) 161

Notes

2D NMR Analysis of the polylactone derivative of colominic acid. Complete ^{1}H and ^{13}C NMR chemical shift assignments
T.M. Flaherty and J. Gervay (Tucson, AZ, USA) 173

An improved synthesis of 1,6-anhydro-2,3-di-*O*-benzyl-β-D-*xylo*-hexopyranos-4-ulose
S. Caron, A.I. McDonald and C.H. Heathcock (Berkeley, CA, USA) 179

Preparation of D-sorbose from L-glucitol by bioconversion with *Pseudomonas* sp. Ac
A. Huwig, S. Emmel and F. Giffhorn (Saarbrücken, Germany) 183

Corrigendum C1

Book review C3

Announcement C5

Information for authors C7

Regular papers

Analysis of the structural heterogeneity of laminarin by electrospray-ionisation–mass spectrometry
S.M. Read, G. Currie and A. Bacic (Parkville, Australia) 187

Acid modification of starch granules in alcohols: effects of temperature, acid concentration, and starch concentration
J.F. Robyt, J.-y. Choe, R.S. Hahn and E.B. Fuchs (Ames, IA, USA) 203

Mass spectrometric studies of the thermal decomposition of carbohydrates using ^{13}C-labeled cellulose and glucose
R.J. Evans, D. Wang, F.A. Agblevor, H.L. Chum (Golden, CO, USA) and S.D. Baldwin (Richmond, VA, USA) 219

Total synthesis of VIM-2 ganglioside isolated from human chronic myelogenous leukemia cells
T. Ehara, A. Kameyama, Y. Yamada (Yokohama, Japan), H. Ishida, M. Kiso and A. Hasegawa (Gifu, Japan) 237

L-Iduronic acid derivatives as glycosyl donors
C. Tabeur, F. Machetto, J.-M. Mallet (Paris, France), P. Duchaussoy, M. Petitou (Toulouse, France) and P. Sinaÿ (Paris, France) 253

Synthesis of (+)-3,7,8-trideoxy-3,7-imino-D-*threo*-L-*galacto*-octitol and its inhibition of β-glucosidases
A. Baudat, S. Picasso and P. Vogel (Lausanne-Dorigny, Switzerland) 277

Structure of a galactan from cell walls of *Bifidobacterium catenulatum* YIT4016
M. Nagaoka, S. Hashimoto, H. Shibata, I. Kimura, K. Kimura, H. Sawada and T. Yokokura (Tokyo, Japan) 285

Notes
The molecular and crystal structure of two trichloroethylidene acetals of epimerized monosaccharides
H. Reinke, D. Rentsch and R. Miethchen (Rostock, Germany) 293

Crystal and molecular structure of 1,6:3,4-dianhydro-2-*O*-*p*-tolylsulfonyl-β-D-galactopyranose
A.-L. Wang, S.-J. Lu, H.-X. Fu, H.-Q. Wang (Lanzhou, P.R. China) and X.-Y. Huang (Fuzhou, P.R. China) 301

Synthesis of 3-acetamido-1,2,4-tri-*O*-acetyl-3,5-dideoxy-5-*C*-[(*R*)- and (*S*)-phenylphosphinyl]-α- and -β-D-xylopyranoses
K. Seo (Numazu, Japan), M. Yamashita, T. Oshikawa (Hamamatsu, Japan) and J.-i. Kobayashi (Fukuroi, Japan) 307

Synthesis of 2-acetamido-3-*O*-acetyl-2-deoxy-D-mannose phosphoramidites
S.J. Freese and W.F. Vann (Bethesda, MD, USA) 313

New imino sugar derivatives from a 6-azido-6-deoxy sugar formazan
V. Zsoldos-Mády, I. Pintér, P. Sándor and A. Messmer (Budapest, Hungary) 321

Announcement C9

Author index C11

Subject index C13

Contents C19